FIGURE 8. $\sigma(\mathbf{p})$: When $c(\mathbf{p})$ is an edge.

Theorem 5. Given $\mathbf{o}^v \in \mathcal{O}_n^+$, the cardinality of $[\mathbf{o}^v]^+$ is

$$(4.3) \quad |[\mathbf{o}^v]^+| = \frac{2\epsilon(\mathbf{p})}{\sigma(\mathbf{p})},$$

where $\mathbf{p} = U(d(\mathbf{o}^v))$, and $\epsilon(\mathbf{p})$ is the number of edges in \mathbf{p} .

Proof. Since $\bar{d}(RLo^v) = \bar{d}(\mathbf{o}^v)$ and $U(d(RLo^v)) = U(d(\mathbf{o}^v))$, the size of $[\mathbf{o}^v]^+$ is determined by \mathbf{p} . More precisely, the size of $[\mathbf{o}^v]^+$ equals the number of ways of identifying a vertex in $w \in \mathbf{p}$ with $v \in \bar{d}(\mathbf{o}^v)$. For each vertex w in \mathbf{p} , since \mathbf{p} is embedded in the plane, we have $\deg(w)$ distinct ways of identifying w with $v \in d(\mathbf{o}^v)$. So, if we allow repetition, the number of all possible ways of attaching \mathbf{p} to $\bar{d}(\mathbf{o}^v)$ is $\sum_{w \in \mathbf{p}} \deg(w) = 2\epsilon(\mathbf{p})$. But by the definition of the symmetry number, each pattern occurs exactly $\sigma(\mathbf{p})$ times. This yields (4.3). \square

REFERENCES

- [1] F. Bergeron, G. Labelle and P. Leroux, *Combinatorial Species and Tree-like Structures*, Encyclopedia of Mathematics and its Applications vol. 67, Cambridge University Press, 1998.
- [2] C. Chauve, *Half of the nodes of Catalan trees are leaves*, personal communication.
- [3] B. Dasarathy and C. Yang, A transformation on ordered trees, *Comput. J.*, **23**, No. 2 (1980), 161–164.
- [4] N. Dershowitz and S. Zaks, Applied tree enumerations, in *Lecture Notes in Computer Science*, vol. 112, 180–193, Springer-Verlag, Berlin, 1981.
- [5] E. Deutsch, An involution on Dyck paths and its consequences, *Discrete Math.*, **204** (1999), 163–166.
- [6] F. Harary and E. Palmer, *Graphical Enumeration*, Academic Press, New York, 1973.
- [7] D. E. Knuth, *The Art of Computer Programming*, vol. 1, Fundamental Algorithms, 2nd ed., Addison-Wesley, Reading, Massachusetts, 1973.
- [8] S. Seo, A pairing of vertices in ordered trees, *Discrete Math.*, **241** (2001), 471–477.
- [9] N. J. A. Sloane, *On-Line Encyclopedia of Integer Sequences*, <http://www.research.att.com/~njas/sequences/>.

DEPARTMENT OF MATHEMATICS, KOREA ADVANCED INSTITUTE OF SCIENCE AND TECHNOLOGY, TAEJON, 305-701, REPUBLIC OF KOREA

E-mail address: shseo@math.kaist.ac.kr, chattest@math.kaist.ac.kr

PERIODIC DE BRUIJN TRIANGLES: EXACT AND ASYMPTOTIC RESULTS

B. SHAPIRO, M. SHAPIRO, AND A. VAINSHTEIN

ABSTRACT. We study the distribution of the number of permutations with a given periodic up-down sequence w.r.t. the last entry, find exponential generating functions and prove asymptotic formulas for this distribution.

RÉSUMÉ. Nous étudions la distribution du nombre de permutations ayant une suite périodique donnée de montées-descentes par rapport leur dernière entrée. Nous en trouvons les séries génératrices et montrons des formules asymptotiques pour cette distribution.

Let $\sigma = (\sigma_1, \dots, \sigma_n)$ be a permutation of length n . We associate with σ its *up-down sequence* (sometimes called the *shape* of σ , or the *signature* of σ) $\mathcal{P}(\sigma) = (p_1, \dots, p_{n-1})$, which is a binary vector of length $n - 1$ such that $p_i = 1$ if $\sigma_i < \sigma_{i+1}$ and $p_i = 0$ otherwise. During the last 120 years, many authors have studied the number $\sharp_n^{\mathcal{P}}$ of all permutations of length n with a given up-down sequence \mathcal{P} . Apparently, for the first time this problem was investigated by D. André [An1, An2], who considered the so-called alternating (or up-down) permutations corresponding to the sequence $\mathcal{P} = (1, 0, 1, 0, \dots) = (10)^*$ and proved that the exponential generating function for the number of such permutations is equal to $\tan x + \sec x$. In [An3] he proved that this number grows asymptotically as $2n!(2/\pi)^{n+1}$.

A general approach to this problem was suggested by MacMahon (see [MM]). This approach leads to determinantal formulas for $\sharp_n^{\mathcal{P}}$, rediscovered later by Niven [Ni] from very basic combinatorial considerations. For the relations of this approach to the representation theory of the symmetric group, and for its generalizations, see [Fo, St, BW].

Another, purely combinatorial approach to the same problem was suggested by Carlitz [Ca11]. His general recursive formula for $\sharp_n^{\mathcal{P}}$ is rather difficult to use. However, he managed to obtain explicit expressions for the corresponding exponential generating functions for certain *periodic* cases, that is for up-down sequences of the form $\mathcal{P} = (p)^*$, where p is a binary vector of a fixed length called the *period* of \mathcal{P} . In [Ca11, Ca2] he considered the case $\mathcal{P} = (1^k 0)^*$ and expressed the corresponding generating function via the *Olivier functions of the k th order*

$$\varphi_{k,i}(x) = \sum_{j=0}^{\infty} \frac{x^{jk+i}}{(jk+i)!}, \quad 0 \leq i \leq k-1.$$

Another case, $\mathcal{P} = (1^2 0^2)^*$, was considered in [CS1, CS2] and solved via Olivier functions of the fourth order. It follows that asymptotically $\sharp_n^{\mathcal{P}}$ in this case grows as $4n!(2/\gamma)^{n+1}$, where $\gamma = 3.7502\dots$ is the smallest positive solution of the equation $\cos t \cosh t + 1 = 0$.

The general periodic problem was solved completely in [CGJN]. As in the two particular cases mentioned above, the answer is expressed via Olivier functions. The techniques used involves matrix Riccati equations, and is rather complicated.

An additional dimension in the problem was introduced by Entringer [En] who studied the distribution of the alternating permutations by the last entry. He observed that the number $\sharp_{i,j}$ of alternating permutations of length i whose last entry equals j satisfy the

following equations:

$$\begin{aligned} \#_{i,j} &= \#_{i,j-1} + \#_{i-1,j-1}, & \#_{i,1} &= 0, & i &= 2k, k > 0, \\ \#_{i,j} &= \#_{i,j+1} + \#_{i-1,j}, & \#_{i,i} &= 0, & i &= 2k + 1, k > 0, \end{aligned} \tag{1}$$

with $\#_{1,1} = 1$. These equations can be represented graphically as the following triangle displayed on Figure 1.




FIGURE 1. The Entringer triangle

Each even row of the triangle starts with 0, and an entry in such a row is equal to the sum of its *left* neighbors in the current and in the previous rows. Similarly, each odd row (except for the first one) ends with 0, and an entry in such a row is equal to the sum of its *right* neighbors in the current and in the previous rows.

The Entringer triangle was studied by many authors. In particular, Arnold [Ar1, Ar2] gave an interpretation of the entries of this triangle in terms of real polynomials with real critical values. Besides, he considered the exponential generating function

$$A(x, y) = \sum_{i \geq 1} \sum_{j=1}^i (-1)^{i(i-1)/2} \#_{i,j} \frac{x^{i-j} y^{j-1}}{(i-j)!(j-1)!}$$

and proved that $A(x, y) = e^y / \cosh(x + y)$. In fact, $A(x, y)$ is the generating function of the *signed Entringer triangle*, which is obtained from the ordinary one by reversing signs in each i th row, where i equals 2 or 3 modulo 4. Observe that the entries $\tilde{\#}_{i,j}$ of the signed Entringer triangle satisfy relations

$$\tilde{\#}_{i,j} = \tilde{\#}_{i,j-1} + \tilde{\#}_{i-1,j-1} \tag{2}$$

with boundary conditions $\tilde{\#}_{i,1} = 0$ for $i = 2k$, $\tilde{\#}_{i,i} = 0$ for $i = 2k + 1$, $k > 0$, $\tilde{\#}_{1,1} = 1$. General triangles satisfying relation (2) with arbitrary boundary conditions were first studied more than 120 years ago by Seidel [Se]. In particular, he proved that the ratio of exponential generating functions for the numbers on the right and on the left sides of such a triangle equals e^x . More recently such triangles were studied, from the combinatorial point of view, in [DV, Du1, Du2]. In particular, it is proved in [DV] that the exponential generating function for a Seidel triangle is equal to $e^y F(x+y)$, where $F(x)$ is the corresponding function for the left side of the triangle.

The case of general up-down sequences was addressed by de Bruijn [dB] (see also [Vi] for another version of the same result). Let $\#_{i,j}^{\mathcal{P}}$ be the number of permutations of length i whose last entry equals j and whose up-down sequence equals $\mathcal{P} = (p_1, p_2, \dots)$. He proved

that these numbers satisfy the following equations:

$$\begin{aligned} \#_{i,j}^{\mathcal{P}} &= \#_{i,j-1}^{\mathcal{P}} + \#_{i-1,j-1}^{\mathcal{P}}, & \#_{i,1}^{\mathcal{P}} &= 0, & \text{if } p_{i-1} &= 1 \\ \#_{i,j}^{\mathcal{P}} &= \#_{i,j+1}^{\mathcal{P}} + \#_{i-1,j}^{\mathcal{P}}, & \#_{i,i}^{\mathcal{P}} &= 0, & \text{if } p_{i-1} &= 0. \end{aligned}$$

with $\#_{1,1}^{\mathcal{P}} = 1$. Evidently, for $\mathcal{P} = (10)^*$ one gets the Entringer relations (1). As before, these equations can be represented graphically as a triangle, and the direction in which one has to advance along the rows of the triangle is governed by the sequence \mathcal{P} . We call this triangle the *de Bruijn triangle* corresponding to the up-down sequence \mathcal{P} . A de Bruijn triangle is said to be *periodic* if the corresponding up-down sequence is periodic.

The *signed de Bruijn triangle* is obtained from the ordinary de Bruijn triangle by multiplying its i th row by $\varepsilon_i = (-1)^{p_1+p_2+\dots+p_{i-1}+i-1}$, $i \geq 2$. The corresponding exponential generating function is defined by

$$F^{\mathcal{P}}(x, y) = \sum_{i \geq 1} \sum_{j=1}^i \varepsilon_i \#_{i,j}^{\mathcal{P}} \frac{x^{i-j} y^{j-1}}{(i-j)!(j-1)!}.$$

Let \mathcal{P} be a periodic up-down sequence with period p of length m , and let $i_1 < i_2 < \dots < i_r$ be the locations of zeros in p . Without loss of generality we assume that $i_r = m$ (otherwise we consider instead of \mathcal{P} the up-down sequence $\bar{\mathcal{P}} = (\bar{p})^*$, where $\bar{p}_i = 1 - p_i$ for $1 \leq i \leq m$; evidently, the de Bruijn triangle for $\bar{\mathcal{P}}$ is obtained from that for \mathcal{P} by the reflection in the main diagonal).

Theorem 1. *The exponential generating function of the signed periodic de Bruijn triangle corresponding to the up-down sequence \mathcal{P} is given by $F^{\mathcal{P}}(x, y) = e^y f^{\mathcal{P}}(x + y)$, where*

$$f^{\mathcal{P}}(t) = \frac{\det \bar{M}^{\mathcal{P}}(t)}{\det M^{\mathcal{P}}(t)}$$

and $M^{\mathcal{P}}(t)$ and $\bar{M}^{\mathcal{P}}(t)$ are $r \times r$ matrices

$$M^{\mathcal{P}}(t) = \begin{pmatrix} \varphi_{m,0} & t^{i_1} \varphi_{m,n-i_1} & t^{i_2} \varphi_{m,n-i_2} & \dots & t^{i_{r-1}} \varphi_{m,n-i_{r-1}} \\ t^{-i_1} \varphi_{m,i_1} & \varphi_{m,0} & t^{i_1} \varphi_{m,n-i_1} & \dots & t^{i_{r-2}} \varphi_{m,n-i_{r-2}} \\ t^{-i_2} \varphi_{m,i_2} & t^{-i_1} \varphi_{m,i_1} & \varphi_{m,0} & \dots & t^{i_{r-3}} \varphi_{m,n-i_{r-3}} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ t^{-i_{r-1}} \varphi_{m,i_{r-1}} & t^{-i_{r-2}} \varphi_{m,i_{r-2}} & t^{-i_{r-3}} \varphi_{m,i_{r-3}} & \dots & \varphi_{m,0} \end{pmatrix}$$

and

$$\bar{M}^{\mathcal{P}}(t) = \begin{pmatrix} 1 & t^{i_1} & t^{i_2} & \dots & t^{i_{r-1}} \\ t^{-i_1} \varphi_{m,i_1} & \varphi_{m,0} & t^{i_1} \varphi_{m,n-i_1} & \dots & t^{i_{r-2}} \varphi_{m,n-i_{r-2}} \\ t^{-i_2} \varphi_{m,i_2} & t^{-i_1} \varphi_{m,i_1} & \varphi_{m,0} & \dots & t^{i_{r-3}} \varphi_{m,n-i_{r-3}} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ t^{-i_{r-1}} \varphi_{m,i_{r-1}} & t^{-i_{r-2}} \varphi_{m,i_{r-2}} & t^{-i_{r-3}} \varphi_{m,i_{r-3}} & \dots & \varphi_{m,0} \end{pmatrix}$$

with $\varphi_{m,j} = \varphi_{m,j}(t)$.

The proof of the theorem follows easily from the above mentioned properties of Seidel triangles and an evident identity $e^x = \sum_{i=0}^{m-1} \varphi_{m,i}(x)$. As a corollary we get the result of [CGJN] cited above.

Moreover, the same techniques allows to obtain generating functions for other Seidel triangles with periodic boundary conditions, such as the triangle for Genocchi numbers (see [DV]). It can be also extended to pairs of Seidel triangles with periodic boundary conditions, such as Arnold triangles $L(\beta)$ and $R(\beta)$ for Springer numbers (see [Ar2, Du2]), thus recovering several combinatorial results obtained in [Sp, Ar2].

Let us now consider the asymptotic behavior of the numbers $\sharp_{i,j}^{\mathcal{P}}$. It was observed without a proof in [Ar2, p. 18] that the even rows of the ordinary Entringer triangle approximate, after an appropriate normalization, the function $\sin x$ on the interval $[0, \pi/2]$, while the odd rows approximate $\cos x$. Exact statements with the first two correction terms can be found in [SY].

We generalize this result to arbitrary periodic de Bruijn triangles. Consider the space \mathcal{L}_2 of all L_2 -functions on the interval $[0, \pi/2]$. Let \mathfrak{S}_k be the linear space of all sequences of real numbers of length k . We define the standard inclusion $i_k \mathfrak{S}_k \hookrightarrow \mathcal{L}_2$ as the linear map sending a sequence $\{a_1, \dots, a_k\}$ to the function whose value equals a_j on the interval $[\frac{\pi(j-1)}{2k}, \frac{\pi j}{2m})$.

For any $l, 0 \leq l \leq m - 1$, let $\mathcal{N}_{k,l}^{\mathcal{P}}$ denote the sequence $\{\sharp_{kn+l,j}^{\mathcal{P}}\} \in \mathfrak{S}_{km+l}$, and let $\xi_{k,l}^{\mathcal{P}} = ci_{km+l}(\mathcal{N}_{k,l}^{\mathcal{P}})$, where c is a normalizing coefficient depending on k and l and chosen in such a way that $\int_0^{\pi/2} \xi_{k,l}^{\mathcal{P}}(\theta) d\theta = 1$.

Theorem 2. *For any $l, 0 \leq k \leq m - 1$, the sequence of functions $\xi_{k,l}^{\mathcal{P}}$ converges in the L_2 -metric to the normalized first eigenfunction $\Xi_l^{\mathcal{P}}$ of the two-point spectral problem*

$$z^{(m)} = (-1)^r \lambda^m z$$

with m homogeneous boundary conditions

$$\begin{aligned} z^{(i)}(0) &= 0 && \text{if } pl+i-1 = 1, \\ z^{(i)}(\pi/2) &= 0 && \text{if } pl+i-1 = 0, \end{aligned}$$

where r is the number of zeros in the period p .

The first eigenvalue of the above spectral problem is the minimal positive solution of the equation

$$\det \tilde{M}^{\mathcal{P}}(\varepsilon \lambda \pi/2) = 0,$$

where ε is the m th primitive root of $(-1)^r$ and

$$\tilde{M}^{\mathcal{P}}(t) = \begin{pmatrix} \varphi_{m,0} & \varphi_{m,i_2-i_1} & \varphi_{m,i_3-i_1} & \cdots & \varphi_{m,i_r-i_1} \\ \varphi_{m,i_1-i_2} & \varphi_{m,0} & \varphi_{m,i_3-i_2} & \cdots & \varphi_{m,i_r-i_2} \\ \varphi_{m,i_1-i_3} & \varphi_{m,i_2-i_3} & \varphi_{m,0} & \cdots & \varphi_{m,i_r-i_3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \varphi_{m,i_1-i_r} & \varphi_{m,i_2-i_r} & \varphi_{m,i_3-i_r} & \cdots & \varphi_{m,0} \end{pmatrix}$$

with $\varphi_{m,j} = \varphi_{m,j}(t)$.

The eigenfunction $\Xi_l^{\mathcal{P}}$ is normalized by the condition $\int_0^{\pi/2} \Xi_l^{\mathcal{P}}(\theta) d\theta = 1$, and the indices are understood modulo m .

In particular, for the Entringer numbers $\sharp_{i,j}$ one gets the sine law of [SY]:

$$\begin{cases} \lim_{k \rightarrow \infty, \frac{j}{2k+1} \rightarrow u} \sharp_{2k+1,j} = 2(2k)! \left(\frac{2}{\pi}\right)^{2k+1} \cos \frac{\pi u}{2}, \\ \lim_{k \rightarrow \infty, \frac{j}{2k} \rightarrow u} \sharp_{2k,j} = 2(2k-1)! \left(\frac{2}{\pi}\right)^{2l} \sin \frac{\pi u}{2}. \end{cases}$$

The starting point of this research was the result by the first and the third author that the numbers $\sharp_{i,j}^{\mathcal{P}}$ for $\mathcal{P} = (1^2 0^2)^*$ arise naturally in counting real rational functions of a certain type, see [SV]. The authors are grateful to Max-Planck-Institut für Mathematik, Bonn for its hospitality in September 2000 and to the Royal Institute of Technology, Stockholm for the financial support of the visit of A. V. to Stockholm in July-August 2001. Sincere thanks goes to Prof. Harold Shapiro from The Royal Institute of Technology for his help with functional analysis in the proof of Theorem 2.

REFERENCES

- [An1] D. André, Développments de $\sec x$ et de $\tan x$, *C.R. Heb.* **88** (1879), 965–967.
- [An2] D. André, Sur les permutations alternées, *J. Math. Pures Appl.* **7** (1881), 167–184.
- [An3] D. André, Probabilité pour qu’une permutation donnée de n lettres soit une permutation alternée, *C.R. Heb.* **97** (1883), 983–984.
- [Ar1] V. Arnold, Bernoulli–Euler updown numbers associated with functions singularities, their combinatorics and arithmetics, *Duke Math. J.* **63** (1991), 537–555.
- [Ar2] V. Arnold, The calculus of snakes and the combinatorics of Bernoulli, Euler and Springer numbers of Coxeter groups, *Russ. Math. Surv.* **47** (1992), 1–51.
- [BW] A. Björner and M. Wachs, Generalized quotients in Coxeter groups, *Trans. Amer. Math. Soc.* **308** (1988), 1–37.
- [Ca1] L. Carlitz, Permutations with prescribed pattern, *Math. Nachr.* **58** (1973), 31–53.
- [Ca2] L. Carlitz, Generating functions for a special class of permutations, *Proc. Amer. Math. Soc.* **47** (1975), 251–256.
- [CS1] L. Carlitz and R. Scoville, Enumeration of rises and falls by position, *Discr. Math.* **5** (1973), 45–59.
- [CS2] L. Carlitz and R. Scoville, Generating functions for certain types of permutations, *J. Comb. Theory A* **18** (1975), 262–275.
- [CGJN] C. Collins, I. Goulden, D. Jackson, and O. Nierstrasz, A combinatorial application of matrix Riccati equations and their q -analogue, *Discr. Math.* **36** (1981), 139–153.
- [dB] N. de Bruijn, Permutations with given ups and downs, *Nieuw Arch. Wisk.* **18** (1970), 61–65.
- [Du1] D. Dumont, Matrices d’Euler–Seidel, *Sem. Lothar. Comb.* **5** (1981), B05c.
- [Du2] D. Dumont, Further triangles of Seidel–Arnold type and continued fractions related to Euler and Springer numbers, *Adv. Appl. Math.* **16** (1995), 275–296.
- [DV] D. Dumont and G. Viennot, A combinatorial interpretation of the Seidel generation of Genocchi numbers, *Ann. Discr. Math.* **6** (1980), 77–87.
- [En] R. Entringer, A combinatorial interpretation of the Euler and Bernoulli numbers, *Nieuw Arch. Wisk.* **14** (1966), 241–246.
- [Fo] H. Foulkes, Enumeration of permutations with prescribed up-down and inversion sequences, *Discr. Math.* **15** (1976), 235–252.
- [MM] P. MacMahon, Second memoir on the compositions of numbers, *Philos. Trans. Roy. Soc. London (A)* **207** (1908), 65–134.
- [Ni] I. Niven, A combinatorial problem of finite sequences, *Nieuw Arch. Wisk.* **16** (1968), 116–123.
- [Se] L. Seidel, Über eine einfache Entstehungsweise der Bernoullischen Zahlen und einiger verwandten Reihen, *Sitzungsber. Münch. Akad.* **4** (1877), 157–187.
- [Sp] T. Springer, Remarks on a combinatorial problem, *Nieuw Arch. Wisk.* **19** (1971), 30–36.
- [St] R. Stanley, Binomial posets, Möbius inversion, and permutation enumeration, *J. Comb. Theory A* **20** (1976), 336–356.
- [SY] G. Simons and Y-C. Yao, Asymptotic Sine laws arising from alternating random permutations and sequences, *Random Structures and Algorithms* **8** (1996), 75–95.
- [SV] B. Shapiro and A. Vainshtein, Counting real rational functions with all real critical values, (in preparation).
- [Vi] G. Viennot, Permutations ayant une forme donnee, *Discrete Math.* **26** (1979), 279–284.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF STOCKHOLM, S-10691, SWEDEN
E-mail address: shapiro@matematik.su.se

DEPARTMENT OF MATHEMATICS, MICHIGAN STATE UNIVERSITY, EAST LANSING, MI 48824, USA
E-mail address: mshapiro@math.msu.edu

DEPARTMENT OF MATHEMATICS AND DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF HAIFA,
 MOUNT CARMEL, 31905 HAIFA, ISRAEL
E-mail address: alek@cs.haifa.ac.il