

MULTICOMPLEXES AND POLYNOMIALS WITH REAL ZEROS.

JASON BELL AND MARK SKANDERA

ABSTRACT. We show that each polynomial $a(z) = 1 + a_1z + \cdots + a_dz^d$ in $\mathbb{N}[z]$ having only real zeros is the f -polynomial of a multicomplex. Evidently $a(z)$ is also the h -polynomial of a Cohen-Macaulay ring and is the g -polynomial of a simplicial polytope.

RÉSUMÉ. Nous montrons que chaque polynôme $a(z) = 1 + a_1z + \cdots + a_dz^d$ dans $\mathbb{N}[z]$ qui ne possède que des zéros réelles c'est le polynôme- f d'un multicomplexe. Évidemment $a(z)$ est aussi le polynôme- h d'un anneau de Cohen-Macaulay et est le polynôme- g d'un polytope simplicial.

1. INTRODUCTION

Several interesting results and open questions in algebraic combinatorics concern simplicial complexes and polynomials

$$(1.1) \quad a(z) = 1 + a_1z + \cdots + a_dz^d \in \mathbb{N}[z]$$

having only real zeros. A few examples are the following.

1. The f -polynomial of a $(\mathbf{3} + \mathbf{1})$ -free poset has only real zeros [5], [14], [17].
2. The f -polynomial of a matching complex has only real zeros [8].
3. The f -polynomial of a distributive lattice is conjectured to have only real zeros [12].
4. If P is a series-parallel poset, then the f -polynomial of the distributive lattice $J(P)$ has only real zeros [19].
5. The question of whether the f -polynomial of a modular lattice has only real zeros is open [18].

Progress on the open questions above and on related open questions is obstructed somewhat by the lack of a known combinatorial interpretation for the coefficients of a polynomial (1.1) having only real zeros. A particularly nice combinatorial interpretation might involve faces in a simplicial complex.

Question 1.1. Let the polynomial $a(z) = 1 + a_1z + \cdots + a_dz^d$ have nonnegative integer coefficients and only real zeros. Is $a(z)$ the f -polynomial of a simplicial complex?

The more general class of multicomplexes might also provide a combinatorial interpretation. In Section 2 we will define the f -polynomials of simplicial complexes and multicomplexes, and we will summarize the well-known characterizations of these

polynomials. In Section 3 we will state inequalities satisfied by the coefficients of polynomials with real zeros. These inequalities lead to a proof in Section 4 that each polynomial (1.1) having only real zeros is the f -polynomial of a multicomplex.

2. CHARACTERIZATION OF THE f -VECTORS OF MULTICOMPLEXES AND SIMPLICIAL COMPLEXES

A *multicomplex* on a set $\{x_1, \dots, x_n\}$ of variables is a collection Σ of monomials in x_1, \dots, x_n which satisfies

1. The monomial x_i belongs to Σ , for $i = 1, \dots, n$.
2. If the monomial u belongs to Σ and w divides u , then w also belongs to Σ .

A multicomplex Σ is called a *simplicial complex* if each monomial in Σ is square-free.

Let Σ be a multicomplex on x_1, \dots, x_n . We define the f -vector of Σ to be the sequence

$$(2.1) \quad f_\Sigma = (a_i)_{i \geq 0},$$

where a_i is the number of monomials of degree i in Σ . Note that we necessarily have $a_0 = 1$, unless $n = 0$. Also note that the f -vector of a simplicial complex has only finitely many nonzero components.

Two well-known theorems characterize the f -vectors of multicomplexes and simplicial complexes in terms of functions based upon the following expression of a positive integer m as a sum of binomial coefficients. Given a positive integer i , we define the i th *Macaulay expansion* of m to be the unique expression

$$(2.2) \quad m = \binom{n_i}{i} + \binom{n_{i-1}}{i-1} + \dots + \binom{n_j}{j},$$

satisfying

$$n_i > n_{i-1} > \dots > n_j \geq j \geq 1.$$

To obtain this expression we choose n_i to be the unique positive integer which satisfies

$$\binom{n_i}{i} \leq m < \binom{n_i + 1}{i},$$

and then we compute the $(i-1)$ st Macaulay expansion of $m - \binom{n_i}{i}$.

We define the families $(\mu_i)_{i \geq 1}$, $(\kappa_i)_{i \geq 1}$ of functions on \mathbb{N} by

$$\mu_i(m) = \begin{cases} \binom{n_i+1}{i+1} + \binom{n_{i-1}+1}{i} + \dots + \binom{n_j+1}{j+1} & \text{if } m > 0, \\ 0 & \text{otherwise.} \end{cases}$$

$$\kappa_i(m) = \begin{cases} \binom{n_i}{i+1} + \binom{n_{i-1}}{i} + \dots + \binom{n_j}{j+1} & \text{if } m > 0, \\ 0 & \text{otherwise.} \end{cases}$$

The characterization of f -vectors of multicomplexes is due to Macaulay [11].

Theorem 2.1. *An integer sequence (a_0, a_1, \dots) is the f -vector of a nonempty multicomplex on n variables if and only if we have $a_0 = 1$, $a_1 = n$ and*

$$0 \leq a_{i+1} \leq \mu_i(a_i)$$

for $i \geq 1$.

The characterization of f -vectors of simplicial complexes is due (independently) to Kruskal [10], Katona [9], and Schützenberger [13].

Theorem 2.2. *An integer sequence (a_0, \dots, a_d) is the f -vector of a nonempty simplicial complex on n variables if and only if we have $a_0 = 1$, $a_1 = n$ and*

$$0 < a_{i+1} \leq \kappa_i(a_i)$$

for $i = 1, \dots, d - 1$.

(See [3], [6] for proofs of these theorems.)

The functions μ_i and κ_i may be expressed in terms of one another very easily.

Proposition 2.3. *For any positive integers m, i , we have*

$$\kappa_i(m) + m = \mu_i(m).$$

Proof. Omitted. □

It is customary to define the f -vector of a finite multicomplex Σ to be only the nonzero subsequence of the sequence (2.1),

$$f_\Sigma = (a_0, \dots, a_d).$$

We then define the f -polynomial of Σ to be

$$f_\Sigma(z) = a_0 + a_1z + \dots + a_dz^d.$$

We may also associate f -vectors and f -polynomials to posets. In particular, the set of chains of a poset P forms a simplicial complex $\mathcal{O}(P)$ called the *order complex* of P . (See [16, Ch. 3].) We then define the f -vector f_P and f -polynomial $f_P(z)$ of P to be $f_{\mathcal{O}(P)}$ and $f_{\mathcal{O}(P)}(z)$, respectively.

Multicomplexes have an important interpretation in commutative algebra: if R is a graded k -algebra generated by elements x_1, \dots, x_n , then R has a k -basis which is a multicomplex on x_1, \dots, x_n . Furthermore, $a(z)$ is the f -polynomial of a finite multicomplex if and only if for any nonnegative integer c there exists a c -dimensional Cohen-Macaulay ring whose Hilbert series is

$$\frac{a(z)}{(1-z)^c}.$$

(See [15, pp. 56-57].)

3. INEQUALITIES PERTAINING TO POLYNOMIALS WITH REAL ZEROS

Let the polynomial $a(z) = 1 + a_1z + \cdots + a_dz^d$ in $\mathbb{R}[z]$ have positive coefficients. Conditions on the sequence $(1, a_1, \dots, a_d)$ which are both necessary and sufficient for $a(z)$ to have only real zeros are known but somewhat cumbersome. (See e.g., [1, Thm. 1], [4, p. 203].) On the other hand, several well-known conditions which are merely necessary are quite simple. In the event that $a(z)$ has only real zeros, the sequence $(1 = a_0, \dots, a_d)$ is *unimodal*,

$$1 \leq \cdots \leq a_j \geq \cdots \geq a_d \quad \text{for some } j,$$

and *log-concave*,

$$a_i^2 \geq a_{i-1}a_{i+1} \quad \text{for } i = 1, \dots, d-1.$$

It also has Newton's log-concavity property,

$$(3.1) \quad \frac{a_i^2}{\binom{d}{i}^2} \geq \frac{a_{i-1}}{\binom{d}{i-1}} \frac{a_{i+1}}{\binom{d}{i+1}} \quad \text{for } i = 1, \dots, d-1,$$

from which one can derive Maclaurin's inequalities [7, p. 52],

$$(3.2) \quad \frac{a_1}{d} \geq \sqrt{\frac{a_2}{\binom{d}{2}}} \geq \sqrt[3]{\frac{a_3}{\binom{d}{3}}} \geq \cdots \geq \sqrt[d]{a_d}.$$

Note that we may interpret (3.2) as a generalization of the Arithmetic Mean - Geometric Mean Inequality by factoring $a(z)$ as $(1 + \beta_1z) \cdots (1 + \beta_dz)$. From Maclaurin's inequalities we obtain the following upper bound for each coefficient a_i in terms of a_1 .

Observation 3.1. *Let $a(z) = 1 + a_1z + \cdots + a_dz^d \in \mathbb{R}[z]$ have positive coefficients and only real zeros. Then for $i = 2, \dots, d$ we have*

$$a_i \leq \binom{d}{i} \left(\frac{a_1}{d}\right)^i.$$

Setting $i = d$ in Observation 3.1 and assuming that all coefficients are integers, we obtain an upper bound on the degree in terms of a_1 .

Observation 3.2. *Let $a(z) = 1 + a_1z + \cdots + a_dz^d \in \mathbb{N}[z]$ have only real zeros. Then d is no greater than a_1 .*

The combination of these two facts yields a third.

Observation 3.3. *For any fixed c there are only finitely many polynomials of the form $1 + cz + a_2z^2 + \cdots + a_dz^d$ in $\mathbb{N}[z]$ which have only real zeros.*

Maclaurin's inequalities also give us a lower bound for each coefficient a_i in terms of a_d . In particular we have the following.

Observation 3.4. Let $a(z) = 1 + a_1z + \cdots + a_dz^d \in \mathbb{N}[z]$ have only real zeros. Then for $i = 1, \dots, d-1$ we have

$$a_i \geq \binom{d}{i}.$$

Thus it is easy to see that a polynomial such as $1 + 4z + 9z^2 + 10z^3 + 5z^4 + z^5$ has nonreal zeros.

A very different consequence of Maclaurin's inequalities relates polynomials with real zeros to the Upper Bound Conjecture for f -vectors of simplicial convex polytopes. (See [15, p. 59] for definitions.)

Proposition 3.5. Let $a(z) = 1 + a_1z + \cdots + a_dz^d \in \mathbb{N}[z]$ have only real zeros and let $f = (f_{-1}, f_0, \dots, f_{d-1})$ be the f -vector of the cyclic polytope $C(a_1, d)$. Then for $i = 1, \dots, d$ we have

$$a_i \leq f_{i-1}.$$

Proof. Define the polynomial

$$b(z) = \left(1 + \frac{a_1}{d}z\right)^d = b_0 + b_1z + \cdots + b_dz^d.$$

By (3.2) a_i is no greater than b_i for $i = 1, \dots, d$. Therefore it suffices to show that b_i is no greater than f_{i-1} for $i = 1, \dots, d$.

By a result of McMullen (see [15, p. 59]), the coefficients of $b(z)$ satisfy the conditions of the Upper Bound Conjecture if the coefficients of the polynomial

$$h(z) = \left(1 + \frac{a_1-d}{d}z\right)^d = 1 + h_1z + \cdots + h_dz^d$$

satisfy

$$h_i \leq \binom{a_1 - d + i - 1}{i}.$$

Computing an upper bound for h_i we have

$$h_i = \binom{d}{i} \left(\frac{a_1 - d}{d}\right)^i = \frac{d(d-1) \cdots (d-i+1)(a_1 - d)^i}{i! d^i} \leq \frac{(a_1 - d)^i}{i!},$$

which is clearly less than or equal to

$$\binom{a_1 - d + i - 1}{i} = \frac{1}{i!} \prod_{j=0}^{i-1} (a_1 - d + j).$$

□

Another consequence of Maclaurin's inequalities is a family of inequalities satisfied by the minors of totally nonnegative matrices. Denote by $\binom{[n]}{k}$ the collection of k -element subsets of $[n] = \{1, \dots, n\}$. For any matrix A of size at least $n \times n$ and any elements S, T of $\binom{[n]}{k}$ define $\Delta_{S,T}$ to be the S, T minor of A , the determinant of the submatrix of A corresponding to rows S and columns T . A matrix is called *totally nonnegative* if all of its minors are nonnegative.

Proposition 3.6. *Let A be an $n \times n$ totally nonnegative matrix and let $k < \ell$ be two integers in $[n]$. Then we have*

$$\binom{n}{k}^\ell \left(\sum_{S \in \binom{[n]}{\ell}} \Delta_{S,S} \right)^k \leq \binom{n}{\ell}^k \left(\sum_{S \in \binom{[n]}{k}} \Delta_{S,S} \right)^\ell.$$

Proof. Suppose A is totally nonnegative. A well-known result states that A has only nonnegative real eigenvalues and therefore that the polynomial

$$\det(Az + I) = 1 + a_1z + \cdots + a_nz^n$$

has only negative real zeros. Since these coefficients are given by

$$a_i = \sum_{S \in \binom{[n]}{i}} \Delta_{S,S},$$

we may apply (3.2) to obtain the desired result. \square

4. MAIN RESULT

In comparing the functions obtained in Section 3 with those described in Section 2, it will be convenient to consider the expression

$$\binom{t}{i} = \frac{t(t-1) \cdots (t-i+1)}{i!}$$

to be a function of a real variable t for any nonnegative integer i . In particular we shall use $\binom{t}{i}$ to define the following function.

Lemma 4.1. *Let i be a positive integer. The real function*

$$\frac{\binom{t}{i+1}}{\binom{t}{i}^{\frac{i+1}{i}}}$$

increases with t on the interval $[i, \infty)$.

Proof. Omitted. \square

Combining Lemma 4.1 with (3.2), we may now relate polynomials with real zeros to f -polynomials of multicomplexes.

Theorem 4.2. *Let $a(z) = 1 + a_1z + \cdots + a_dz^d \in \mathbb{N}[z]$ have only real zeros. Then $a(z)$ is the f -polynomial of a multicomplex.*

Proof. Choose an integer i between 1 and $d-1$. By (3.2) we have

$$(4.1) \quad a_{i+1} \leq \binom{d}{i+1} \left(\frac{a_i}{\binom{d}{i}} \right)^{(i+1)/i}.$$

Now define n_i to be the unique nonnegative integer which satisfies

$$\binom{n_i}{i} \leq a_i < \binom{n_i + 1}{i}.$$

Combining this inequality with Observation 3.4, we obtain

$$\binom{d}{i} \leq a_i < \binom{n_i + 1}{i},$$

which implies that $n_i + 1$ is greater than d . We may therefore apply Lemma 4.1 to replace d by $n_i + 1$ in (4.1),

$$a_{i+1} < \binom{n_i + 1}{i + 1} \left(\frac{a_i}{\binom{n_i + 1}{i}} \right)^{(i+1)/i}.$$

Since a_i is less than $\binom{n_i + 1}{i}$, we also have

$$a_{i+1} < \binom{n_i + 1}{i + 1},$$

which clearly implies that a_{i+1} is less than $\mu_i(a_i)$. \square

Corollary 4.3. *Let $a(z) = 1 + a_1z + \cdots + a_dz^d \in \mathbb{N}[z]$ have only real zeros. Then for any $c \in \mathbb{N}$ there exists a Cohen-Macaulay ring whose Hilbert series is the rational function*

$$\frac{a(z)}{(1 - z)^c}.$$

Equivalently, $a(z)$ is the h -vector of a Cohen-Macaulay complex.

A second consequence of Theorem 4.2 concerns simplicial polytopes. (See [2] for definitions.)

Corollary 4.4. *Let $a(z) = 1 + a_1z + \cdots + a_dz^d \in \mathbb{N}[z]$ have only real zeros. Then for any $c \in \mathbb{N}$ greater than or equal to $2d$, there exists a simplicial c -polytope whose g -polynomial is $a(z)$.*

It would be interesting to strengthen Theorem 4.2 to provide an affirmative answer to Question 1.1. Such an answer seems plausible because if the polynomial

$$(4.2) \quad a(z) = 1 + a_1z + \cdots + a_dz^d \in \mathbb{N}[z]$$

has only real zeros, then the function

$$\binom{d}{i + 1} \left(\frac{a_i}{\binom{d}{i}} \right)^{(i+1)/i}$$

that bounds a_{i+1} is less than $\kappa_i(a_i)$ for a_i large enough. (For instance if a_i is at least $i^2 \binom{d}{i}$.) It follows that for fixed d , at most finitely many polynomials of the form (4.2) have only real zeros and are not f -polynomials of simplicial complexes. It is also possible to show that when $d \leq 4$ or $a_1 \leq 10$, all polynomials of the form (4.2) which have only real zeros are f -polynomials of simplicial complexes.

5. ACKNOWLEDGEMENTS

The authors are grateful to John Stembridge and Joe McClain for their help with computer programming and to Lou Billera, Anders Björner, George Clements, Karen Collins, Mel Hochster, Christopher Smyth, Richard Stanley, Jeff Allotta, Katie Bach, Walter Dulaney, and Eric Hendrickson for many helpful conversations.

REFERENCES

- [1] M. AISSÉN, I. J. SCHOENBERG, AND A. WHITNEY. On generating functions of totally positive sequences. *J. Anal. Math.*, **2** (1952) pp. 93–103.
- [2] L. J. BILLERA AND A. BJÖRNER. Face numbers of polytopes and complexes. In *Handbook of Discrete and Computational Geometry* (J. E. GOODMAN AND J. O’ROURKE, eds.). CRC Press, Boca Raton/New York, 1997 pp. 291–310.
- [3] G. CLEMENTS AND B. LINDSTRÖM. A generalization of a combinatorial theorem of Macaulay. *J. Combin. Theory*, **7** (1969) pp. 230–238.
- [4] F. R. GANTMACHER. *The Theory of Matrices*, vol. 2. Chelsea, New York, 1959.
- [5] V. GASHAROV. Incomparability graphs of $(\mathbf{3} + \mathbf{1})$ -free posets are s -positive. *Discrete Math.*, **157** (1996) pp. 211–215.
- [6] C. GREENE AND D. KLEITMAN. Proof techniques in the theory of finite sets. In *Studies in Combinatorics* (G. C. ROTA, ed.). Mathematical Association of America, 1978 pp. 22–79.
- [7] G. H. HARDY, J. E. LITTLEWOOD, AND G. PÓLYA. *Inequalities*. Cambridge University Press, Cambridge, 1934.
- [8] O. HEILMAN AND E. LIEB. Theory of monomer-dimer systems. *Comm. Math. Physics*, **25** (1972) pp. 190–232.
- [9] G. KATONA. A theorem of finite sets. In *Colloquium on the Theory of Graphs*. Academic Press and Akademiai Kiadó, New York and Budapest, 1968 pp. 187–207.
- [10] J. KRUSKAL. The number of simplices in a complex. In *Mathematical Optimization Techniques*. University of California Press, Berkeley, CA, 1963 pp. 251–278.
- [11] F. S. MACAULAY. Some properties of enumeration in the theory of modular systems. *Proc. London Math. Soc.*, **26** (1927) pp. 531–555.
- [12] J. NEGGERS. Representations of finite partially ordered sets. *J. Combin. Inform. System Sci.*, **3** (1978) pp. 113–133.
- [13] M. P. SCHÜTZENBERGER. A characteristic property of certain polynomials of E. F. Moore and C. E. Shannon. In *RLE Quarterly Progress Report*, 55. MIT Research Laboratory of Electronics, 1959 pp. 117–118.
- [14] M. SKANDERA. A characterization of $(\mathbf{3} + \mathbf{1})$ -free posets. *J. Combin. Theory Ser. A*, **93** (2001) pp. 231–241.
- [15] R. STANLEY. *Combinatorics and Commutative Algebra*. Birkhäuser, Boston, MA, 1996.
- [16] R. STANLEY. *Enumerative Combinatorics*, vol. 1. Cambridge University Press, Cambridge, 1997.
- [17] R. STANLEY. Graph colorings And Related Symmetric functions: Ideas and Applications. *Discrete Math.*, **193** (1998) pp. 267–286.
- [18] R. STANLEY. Positivity problems and conjectures. In *Mathematics: Frontiers and Perspectives* (V. ARNOLD, M. ATIYAH, P. LAX, AND B. MAZUR, eds.). American Mathematical Society, Providence, RI, 2000 pp. 295–319.
- [19] D. WAGNER. Enumeration of functions from posets to chains. *Europ. J. Combin.*, **13** (1992) pp. 313–324.

AUTHORS’ ADDRESS: UNIVERSITY OF MICHIGAN, DEPT. OF MATHEMATICS, 2074 EAST HALL, 525 EAST UNIVERSITY, ANN ARBOR, MI 48109-1109

E-mail address: belljp@umich.edu, mskan@umich.edu